

# Corrigés — Équations différentielles

## Chapitre 7

### Solution 1.

1.  $y(x) = Ce^{2x}$ .
2.  $y(x) = Ce^{-3x}$ .  $y(0) = C = 5$ , donc  $y(x) = 5e^{-3x}$ .
3.  $y(x) = Ce^{-x}$ .  $y(1) = \frac{C}{e} = e$ , donc  $C = e^2$ ,  $y(x) = e^2 e^{-x} = e^{2-x}$ .

### Solution 2.

1.  $y = Ce^x - 2$ .
2.  $y = Ce^{-2x} + 3$ .  $y(0) = C + 3 = 0$ ,  $C = -3$ .  $y(x) = -3e^{-2x} + 3 = 3(1 - e^{-2x})$ .
3.  $y = Ce^{-3x} + 4$ .  $y(0) = C + 4 = 2$ ,  $C = -2$ .  $y(x) = -2e^{-3x} + 4$ .

### Solution 3.

1.  $y = A \cos x + B \sin x$ .
2.  $y(0) = A = 3$ ,  $y'(0) = 3B = 0$ ,  $B = 0$ .  $y = 3 \cos(3x)$ .
3.  $y(0) = A = 0$ ,  $y'(0) = 2B = 2$ ,  $B = 1$ .  $y = \sin(2x)$ .

### Solution 4.

1.  $N(t) = N_0 e^{-\lambda t}$ .
2.  $\frac{N_0}{2} = N_0 e^{-\lambda T} \implies e^{-\lambda T} = \frac{1}{2} \implies T = \frac{\ln 2}{\lambda}$ .

### Solution 5.

$$P(t) = 1000e^{0.03t}. P(t) = 2000 \iff e^{0.03t} = 2 \iff t = \frac{\ln 2}{0.03} \approx 23.1 \text{ ans.}$$

### Solution 6.

$$T(t) = Ce^{-0.05t} + 15. T(0) = C + 15 = 95, C = 80. T(t) = 80e^{-0.05t} + 15. T(20) = 80e^{-1} + 15 \approx 44.4^\circ\text{C}. \text{ Limite : } 15^\circ\text{C (température ambiante).}$$

### Solution 7.

$$(y_1 - y_2)' = y_1' - y_2' = (ay_1 + b) - (ay_2 + b) = a(y_1 - y_2). \text{ Donc résout } z' = az.$$

### Solution 8.

$$y(x) = xe^x + Ce^x = (x + C)e^x.$$

**Solution 9.**

$z = y'$  résout  $z' + 2z = 0$ , donc  $z(x) = Ke^{-2x}$ . Puis  $y(x) = \int z = -\frac{K}{2}e^{-2x} + C = C_1e^{-2x} + C_2$ .